

## Physicians Orders for Donation After Circulatory Death

All orders are to be written by the attending physician or their designee.

No member of the organ recovery team or OPO staff may participate in the guidance or administration of palliative care, or the declaration of death.

### Maintaining Hemodynamics:

- Blood pressure: Systolic >100, MAP > 65-90
- Heart Rate: 60 – 140
- Urine Output goal: 0.5 - 3 ml/kg/hr
- Temperature: 37 - 38° C
- O2% > 95%, PO2 >100

### Nursing:

- NG or OG to LWS
- Turn patient and suction ETT q2H
- Maintain body temp at 96.5-99.5 with warming measures / cooling measures.
- Accurate height and weight

### Labs:

- Lab tubes for serology and HLA collection. These tubes will be provided by the LOOP coordinator.
- Urinalysis Q24 hrs
- 2 Sets of Blood Cultures & a Urine Culture (separate from UA), even if already completed this admission

### **STAT Once, Repeat Q 8 hrs**

- |                                      |  | Order PRN                               | Order Once                                   |
|--------------------------------------|--|---|--|
| <input type="checkbox"/> CMP         | <input type="checkbox"/> Magnesium       | <input type="checkbox"/> ABG Q _____ Hr | <input type="checkbox"/> Blood Type & Screen |
| <input type="checkbox"/> Direct Bili | <input type="checkbox"/> Phosphorus      | <input type="checkbox"/> Troponin       | <input type="checkbox"/> HbgA1C              |
| <input type="checkbox"/> Lactic Acid | <input type="checkbox"/> Ionized Calcium | <input type="checkbox"/> COVID          | <input type="checkbox"/> LDH                 |
| <input type="checkbox"/> PTT         | <input type="checkbox"/> ABG             | <input type="checkbox"/>                | <input type="checkbox"/> GGT                 |
| <input type="checkbox"/> PT / INR    | <input type="checkbox"/> CBC W/ Diff     |   |  |

### Pharmacy:

- MIV: \_\_\_\_\_
- Pressor: \_\_\_\_\_ titrate to keep SBP > 100, MAP > 65.
- Zosyn 3.375g q6H IVPB (or 4.5g q8 depending on hospital formula)
- Electrolyte replacement: Include KCL, Calcium Gluconate, Magnesium and Phosphorus.
- Vasopressin for DI 0.01 U/min (Max 0.05 U/min).

### Additional Orders:

- Insert IV Lines:  Aline,  PIV,  Central Line,  Other: \_\_\_\_\_
- Chest X-ray Q24 hours
- CT Chest, Abdomen, Pelvis
  - W/Contrast, Creat <2.0. Administer 1L LR over 1 hr post CT
  - W/O Contrast, Creat ≥2.0
- Bronchoscopy
  - Sputum Gram Stain
  - Sputum Culture – No AFB
  - COVID BAL Test

### Withdrawal Orders:

- Withdrawal orders and medications per hospital policy to be taken to and/or sent to the operating room for withdrawal.
- Heparin 30,000 units to be administered in the withdrawal location just prior to patient extubation.
- Transfer order if patient does not expire in the allotted time frame for donation.

*Additional orders may be needed. The LOOP coordinator will need to know who to contact for further assistance.*