

Physicians Orders for Donation After Circulatory Death

All orders are to be written by the attending physician or their designee.

No member of the organ recovery team or OPO staff may participate in the guidance or administration of palliative care, or the declaration of death.

Maintaining Hemodynamics:

- Blood pressure: Systolic >100, MAP 60-110
- Heart Rate: 60 – 140
- Urine Output goal: 0.5 - 3 ml/kg/hr
- Temperature: 36.5 - 38° C
- O2% > 95%, PO2 >100

Nursing:

- NG or OG to LWS
- Turn patient and suction ETT q2H
- Maintain body temp at 96.5-99.5 with warming measures / cooling measures.
- Accurate height and weight
- Neuro exam Q12H

Labs:

- Lab tubes for serology and HLA collection. These tubes will be provided by the LOOP coordinator.
- Urinalysis Q24H
- 2 Sets of Blood Cultures with sensitivities & a Urine Culture (separate from UA), even if already completed this admission

STAT Once, Repeat Q 8 hrs

		Order PRN	Order Once
<input type="checkbox"/> CMP	<input type="checkbox"/> Magnesium	<input type="checkbox"/> Troponin	<input type="checkbox"/> Blood Type & Screen
<input type="checkbox"/> Direct Bili	<input type="checkbox"/> Phosphorus	<input type="checkbox"/> CKMB	<input type="checkbox"/> HbgA1C
<input type="checkbox"/> Lactic Acid	<input type="checkbox"/> Ionized Calcium	<input type="checkbox"/> CPK	<input type="checkbox"/> LDH
<input type="checkbox"/> PTT	<input type="checkbox"/> ABG		<input type="checkbox"/> GGT
<input type="checkbox"/> PT / INR	<input type="checkbox"/> CBC W/ Diff		<input type="checkbox"/> COVID
<input type="checkbox"/> ABG			

Pharmacy:

- MIV: _____
- Pressor: _____ titrate to keep SBP > 100, MAP > 60.
- Zosyn 3.375g q6H IVPB (or 4.5g q8H depending on hospital formula)
- Electrolyte replacement: Include KCL, Calcium Gluconate, Magnesium and Phosphorus.
- Vasopressin 0.01-0.04 Units/min to maintain adequate UO

Additional Orders:

- Insert IV Lines: Aline, PIV, Central Line, Other: _____
- Chest X-ray Q24 hours
- CT Chest, Abdomen, Pelvis
 - W/Contrast, Creat < 2.0. Administer 1L LR over 1 hr post CT
 - W/O Contrast, Creat ≥ 2.0
- Bronchoscopy
 - Sputum Gram Stain
 - Sputum Culture – No AFB
 - COVID BAL Test

Withdrawal Orders:

- Withdrawal orders and medications per hospital policy to be taken to and/or sent to the operating room for withdrawal.

- Heparin 30,000 units (**subject to change at the discretion of the hospital care team**) to be provided by the hospital pharmacy and administered in the withdrawal location just prior to patient extubation.
- Transfer order if patient does not expire in the allotted time frame for donation.

Additional orders may be needed. The LOOP coordinator will need to know who to contact for further assistance.